

PROPERTIES

coda-Zn-L is a solution of zinc complexed by lignosulfonates, recommended for foliar or root application.

coda-Zn-L prevents and corrects zinc deficiencies and favors essential enzymatic processes of plants.

It is also a perfect acidifier of the water pH in the nutrient solution, improving its assimilation.

GUARANTEED ANALYSIS		
Zinc (Zn) complexed and water soluble	% w/w	% w/v
	8.0	10.40

Complexing agent: Lignosulfonates.

OBSERVATIONS

coda-Zn-L is compatible with most of the agrochemical products, but a previous compatibility test is recommended.

The mentioned doses are orientative, and may be modified according to climate, crop varieties. Optimal storage temperature is between 5 and 30°C. Not to pile up more than 3 packages or 5 boxes of height.

It is recommended to apply the product under agronomic technical advice.

Causes serious eye damage. May cause damage to organs through prolonged or repeated exposure. Toxic to aquatic life with long lasting effects. Keep out of reach of children. Read label before use. Do not breathe dust, fume, gas, mist, vapors, spray. Avoid release to the environment. Wear protective gloves, protective clothing, eye protection, face protection. If medical advice is needed, have product container or label at hand. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do, continue rinsing. Immediately call a POISON CENTER or doctor/physician. Get medical advice/attention if you feel unwell. Collect spillage. Disposal of waste product and containers must be in accord with local bylaws and regulations.

Contains Zinc Sulphate 10-30%.

24 Hour Emergency Response (NZ):

0800 POISON 0800 764 766

Batch number:

Production date: 04-2024

Expiry date: 04-2027



Sustainable Agro Solutions, S.A.U.
Ctra N-240, Km. 110 - Almacelles - Lleida (Spain)



DANGER

coda-Zn-L



COMPLEXED ZINC SOLUTION

coda

APPLICATION METHOD AND RATES

coda-Zn-L is a product totally soluble in water, indicated for all types of crops, applicable either by foliar or by fertigation.

The recommended average rates are:

Type of application	Application rates
Foliar	200 - 300 mL/100L - application depending on the nutritional status of the crop
Fertigation	3 - 5 L/ha - application 2 - 4 applications per crop cycle

In foliar applications a minimum of 600 liters of water per hectare is recommended in vegetables in general, 1,000 liters of water per hectare in fruit trees and 1,500 liters in citrus fruits.

SHAKE THE PRODUCT BEFORE USING IT
Agricultural Use

Net content: 1 liter



Confirmation of compatibility issued by KIWA BCS and OMRI, compatible product by the regulations **EC no 2021/1165** and **NOP 7 CRF Part 203** for the use of inputs in organic agriculture.

The product is defined and an agricultural compound under the Agricultural Compounds and Veterinary Medicines (ACVM) Act 1997 and is exempt from registration. The product is GHS compliant and is approved under NZ EPA HSR002571 - Fertilisers (Subsidiary Hazard) Group Standard 2020.

New Zealand Importer



67 Elizabeth Avenue, Rakaia 7710
New Zealand **www.sfnz.nz**



SAS, S.A.U. guarantees the quality of the product, regarding the guaranteed analysis stated on this label. The company is not responsible for any damaged caused due to inappropriate usage of the product. Read the label carefully before using the product.

coda